

尼泊尔唐松草 毛茛科一新种

王文采

(中国科学院植物研究所系统与进化植物学国家重点实验室 北京 100093)

摘要 描述了自尼泊尔东部发现的毛茛科唐松草属一新种 尼泊尔唐松草 (*Thalictrum nepalense*)。此新种与叉枝唐松草 (*T. saniculiforme* DC.) 在亲缘关系上甚为接近, 与后者的区别在于新种的茎较高, 有3枚叶和3枝条, 其复单歧聚伞花序的末回分枝极叉开, 其雄蕊花丝上部棍棒状, 下部丝形, 花药在顶端不具细尖头。

关键词 毛茛科 唐松草属 新种 尼泊尔

中图分类号: Q948.15+2 文献标志码: A doi: 10.7525/j.issn.1673-5102.2018.05.001

Thalictrum nepalense a New Species of Ranunculaceae from Nepal

WANG Wen-Tsai

(State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093)

Abstract A species of the genus *Thalictrum* (Ranunculaceae), *T. nepalense*, is described as new from eastern Nepal. It is closely related to *T. saniculiforme* DC., differing from the latter in its taller 3-leaved and 3-branched stem, compound monochasia with only ultimate branches divaricate, above clavate and below filiform stamen filaments, and anthers not apiculate at apex.

Key words Ranunculaceae; *Thalictrum* new species; Nepal

1 大关唐松草 图 1

Thalictrum nepalense W. T. Wang, sp. nov.

Fig. 1.

(Subgen. *Thalictrum* Sect. *Leptostigma* Boivin)

Ob plantas totas glabras, folia 3-ternata, florem carpella numerosa stylis longis apice unciformibus vel circinatis et stigmatibus inconspicuis praedita ferentem species nova haec est arcte affinis *T. saniculiformi* DC., a quo caule altiore 3-foliato et 3-ramoso, monochasiis compositis ter ramosis, tantum eorum ramis ultimis divaricatis, staminum filamentis superne clavatis inferne filiformibus, antheris apice obtusis haud apiculatis differt. In *T. saniculiformi*, caulis humilior, 1-foliatus et 1-ramosus vel nudus, monochasia composita semel vel bis divaricata, staminum fila-

menta anguste linearia, et antherae apice apiculatae sunt.

Perennial herb, totally glabrous. Stem ca. 35 cm tall, above base 2 mm across, 3-leaved and 3-branched, with branches 1-leaved. Basal leaves not seen. Cauline leaves 3-ternate, slightly long or shortly petiolate or sessile; blades 5–8 × 6–8 cm; leaflets papery, obovate, broadly ovate, broadly rhombic or subpentagonal, 0.6–1.7 × 0.4–1.9 cm, at apex rounded, at base broadly cuneate, subcordate or subtruncate, 3-dentate or 3-lobed, with lobes entire or 1–2-dentate, 3–5-nerved, nerves adaxially slightly prominent or subflat, abaxially prominent, with nervules forming conspicuous nervous nets; petiolules filiform, 0.5–8 mm long; petioles 0–5.5 cm long, at base narrowly vaginate. Com-

作者简介: 王文采(1926—)男, 研究员, 中国科学院院士, 从事有花植物分类学研究。

收稿日期: 2018-04-10

Author introduction: WANG Wen-Tsai(1926—) male professor academician of Chinese Academy of Sciences, majoring in taxonomy of flowering plants.

Received date: 2018-04-10




Fig.1 *T. nepalense* A. Roots and basal part of stem ;B. Upper part of flowering plant ;C. Stamen ;D. Carpel ;E. Achene (from holotype)

pound monochasia terminal , including peduncles 5 – 13 cm long , thrice branched and only ultimate branches more or less divaricate ; peduncles 3 – 6.8 cm long ; most bracts foliaceous , or ternate , uppermost ones small , simple , elliptic or obovate , and entire ; pedicels 2 – 5.5 mm long. Flower : Sepals ca-

duous , not seen. Stamens ca. 14 ; filaments 2.5 – 3 mm long , above clavate , as broad as anthers , below filiform ; anthers yellowish , narrow-oblong , ca. 0.8 mm × 0.2 mm , at apex obtuse. Carpels 9 – 12 , sessile ; ovaries ca. 1.2 mm × 0.5 mm ; styles 0.8 – 1.2 mm long , at apex hooked or circinate ; stigmas

inconspicuous. Achenes sessile , compressed , narrowly elliptic , 2. 2 – 3 mm × 0. 25 – 0. 4 mm , on each side longitudinally and thinly 3-ribbed ; persistent styles 0. 8 – 1. 2 mm long , at apex hooked.

E. Nepal : Sagarmatha Zone , Solukhumbu Distr. , Sete(2 500 m). Lamjura La(3 500 m). Taktor (2 850 m) [86° 26′ – 31′ E , 27° 24′ N] 30 July 1997. Coll. M. Wakabayashi , M. Amano , M. Mori , K. R. Rajbhandari & K. Shinozaki No. 9720035 (**holotype** , PE).

In having totally glabrous plants , 3-ternate leaves , flower bearing numerous carpels with long styles hooked or circinate at apex and inconspicuous stigmas this new species is closely related to *T. saniculiforme* DC. also with the characters just stated above , and from the latter differs in its taller 3-leaved and 3-branched stem , thrice branched compound monochasia with only ultimate branches divaricate , above clavate and below filiform filaments of stamens , and not

apiculate anthers. In *T. saniculiforme* , the stems are lower , only 1-leaved and 1-branched , or naked , the compound monochasia are once or twice divaricate , the stamen filaments are narrowly linear , and the anthers are apiculate at apex^[1-3].

Acknowledgements The author is grateful to Mr. Sun Ying-Bao for making the drawing.

References

1. Fu D Z ,Zhu G. Thalictrum[M].//Wu Z Y ,Raven P H. Flora of China. Beijing :Science Press ,St. Louis :Missouri Botanical Garden Press 2001 6 282 – 302.
2. Rau M A. Thalictrum[M].//Sharma B D. Flora of India. Calcutta :Botanical Survey of India ,1993 2 :132 – 143.
3. 王文采. 唐松草属[M].//中国植物志. 北京 :科学出版社 ,1979 27 502 – 592.
Wang W C. Thalictrum[M].//Flora reipublicae popularis sinicae. Beijing Science Press ,1979 27 502 – 592.